MULTIVITAMIN · DISPENSED BY PRESCRIPTION

Folivex

Multivitamin

30 Tablets

Folivex is a prescription multivitamin/multimineral dietary supplement.

- Multivitamin tablets
- 30 Tablets per bottle
- Dispensed by Prescription
- Copay Assistance Program

Visit oncorapharma.com



COUNT	DISPENSING	NDC	FORM
30 Tablets	Dispensed by Prescription	85477-906-30	Tablet

UP TO $75 COPAY ASSISTANCE

Copay Assistance Program

Eligible patients may pay as little as $0 per prescription. The copay card covers up to $75 of a patient's copay.

Not valid for patients enrolled in Medicare, Medicaid, or any other federal or state healthcare program.

Oncora Pharma — COPAY ASSISTANCE CARD

RxBIN: **021940** Group: **99996849**
PCN: **PYS** ID: **91200055846**

Pharmacy Help Desk: 800-800-7364

Supplement Facts (per tablet)

Vitamin A (as Beta-Carotene)	300 mcg RAE	from Folic Acid	0.67 mg DFE
Vitamin C (as Ascorbic Acid)	60 mg	from 5 Methyl Tetrahydrofolate Calcium Salt	1 mg DFE
Vitamin D (as Cholecalciferol)	10 mcg	Vitamin B12 (as Cyanocobalamin)	0.013 mg
Vitamin E (as dl-Alpha Tocopherol Acetate)	4.5 mg (10 IU)	Calcium (as Calcium Carbonate)	80 mg
Vitamin B6 (as Pyridoxine HCl)	26 mg	Magnesium (as Magnesium Oxide)	25 mg
Biotin	0.280 mg	Ferrochel	20 mg
Folate	1.67 mg DFE	Iodine (as Potassium Iodide)	0.150 mg

Other Ingredients: Microcrystalline Cellulose, Maltodextrin, Croscarmellose Sodium, Silicon Dioxide, Stearic Acid, Magnesium Stearate, Film Coating (Hydroxypropyl Methylcellulose, Polyethylene Glycol, Titanium Dioxide, FD&C Blue #1).

Directions

Use only as directed by a healthcare provider. Full Prescribing Information: 833-242-4493 or customerservice@oncorapharma.com.

Storage

Store at 20°–25°C (68°–77°F) excursions permitted to 15°–30°C (59°–86°F) [See USP Controlled Room Temperature]. Avoid excessive heat, light and moisture.

Contraindications

Folivex is contraindicated in patients with a known hypersensitivity to any of the ingredients.

Warnings

Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6.

Keep Out of Reach of Children

Keep this product out of reach of children. In case of accidental overdose, call a doctor or poison control center immediately.

Precautions

Folic acid alone is improper therapy in the treatment of pernicious anemia and other megaloblastic anemias where Vitamin B12 deficient. Folic acid in doses above 1.0 mg daily may obscure pernicious anemia in that hematologic remission can occur while neurological manifestations progress.

Restrictions and eligibility requirements may apply. This program is not insurance. This card is not valid where prohibited by law. See program terms for details. · These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease. · Distributed by: Oncora Pharma, LLC, Dallas, TX 75228.

Oncora Pharma, LLC · 12015 Shiloh Road, Dallas, TX 75228 · oncorapharma.com · customerservice@oncorapharma.com
Oncora Pharma Company Line (toll-free): 833-242-4493
© 2026 Oncora Pharma. All rights reserved.